## IN THE CLAIMS:

Please amend the claims as indicated below:

1. (Currently Amended) A method for transforming gene expression signals, the method comprising the steps of:

determining a plurality of gene expression signals for a gene; and

deriving a transformation <u>for transforming that transforms</u>-the plurality of gene expression signals into transformed gene expression signals for the gene, <u>wherein application of said transformation results resulting</u> in a uniform distribution of the transformed gene expression signals within a selected interval, wherein each gene expression signal is converted by the transformation into a transformed gene expression signal in the selected interval, <u>wherein said transformation allows said transformed gene expression signals to be compared</u>, and wherein the uniform distribution of transformed gene expression signals may be used to determine gene expression patterns.

15

10

5

- 2. (Original) The method of claim 1, further comprising the step of applying the transformation to an additional gene expression signal.
- 3. (Previously presented) The method of claim 1, wherein the step of deriving comprises the steps of:

determining a function that approximates a distribution of the plurality of gene expression signals for the gene; and

using the function to create the transformation.

- 25 4. (Canceled)
  - 5. (Canceled)

	6.	(Canceled)	
	7.	(Canceled)	
5.	8.	(Canceled)	
•	9.	(Canceled)	
10	10.	(Canceled)	
	11.	(Canceled)	
	12.	(Canceled)	
15	13.	(Canceled)	
	14.	(Canceled)	
20	15.	(Canceled)	
20	16.	(Canceled)	
	17.	(Currently Amended) A system comprising:	
25		a memory that stores computer-readable code; and	
25	implement the	a processor operatively coupled to the memory, the processor configured to	
	implement the	lement the computer-readable code, the computer-readable code configured to:  determine a plurality of gene expression signals for a gene; and	
		derive a transformation for transforming that transforms—the plurality of gene	
	expression signals into transformed gene expression signals for the gene, wherein application of		

said transformation results resulting in a uniform distribution of the transformed gene expression signals within a selected interval wherein each gene expression signal is converted by the transformation into a transformed gene expression signal in the selected interval, wherein said transformation allows said transformed gene expression signals to be compared, and wherein the uniform distribution of transformed gene expression signals may be used to determine gene expression patterns.

- 18. (Original) The system of claim 17, wherein the computer-readable code is further configured to apply the transformation to an additional gene expression signal.
- 19. (Previously presented) The system of claim 17, wherein the computer-readable code is further configured, during the step of deriving, to perform the steps of:

determine a function that approximates a distribution of the plurality of gene expression signals for the gene; and

- use the function to create the transformation.
  - 20. (Canceled)

5

10

20

- 21. (Canceled)
- 22. (Canceled)
- 23. (Currently Amended) An article of manufacture comprising:
- a computer readable medium having computer readable code means embodied thereon, the computer readable program code means comprising:
  - a step to determine a plurality of gene expression signals for a gene; and
  - a step to derive a transformation <u>for transforming</u> that transforms the plurality of gene expression signals into transformed gene expression signals for the gene, wherein

application of said transformation results resulting in a uniform distribution of the transformed gene expression signals within a selected interval wherein each gene expression signal is converted by the transformation into a transformed gene expression signal in the selected interval, wherein said transformation allows said transformed gene expression signals to be compared, and wherein the uniform distribution of transformed gene expression signals may be used to determine gene expression patterns.

- 24. (Original) The article of manufacture of claim 23, wherein the computer-readable code means further comprises a step to apply the transformation to an additional gene expression signal.
  - 25. (Previously presented) The article of manufacture of claim 23, wherein the computer-readable code means is further configured, during the step of deriving, to perform:
  - a step to determine a function that approximates a distribution of the plurality of gene expression signals for the gene; and
    - a step to use the function to create the transformation.
  - 26. (Canceled)
- 20 27. (Canceled)

5

15

- 28. (Canceled)
- 29. (Previously presented) The method of claim 1, wherein the selected interval comprises an interval between 0 and 1.